WO 2004/111033

PCT/SE2004/000969

33

Claims

5

1. A compound of formula (I)

$$R^2$$
 N R^3

1

wherein R¹ and R² independently represent phenyl, thienyl or pyridyl each of which is independently optionally substituted by one or more groups represented by Z;

Z represents a C₁₋₈alkyl group, a C₁₋₆alkoxy group, hydroxy, halo, trifluoromethyl, trifluoromethylthio, trifluoromethoxy, trifluoromethylsulphonyl, nitro, mono or di C₁₋₃alkylamido, C₁₋₃alkylsulphonyl, C₁₋₃alkylsulphonyloxy, C₁₋₃alkoxycarbonyl, carboxy, cyano, carbamoyl, mono or di C₁₋₃alkyl carbamoyl, sulphamoyl, acetyl, an aromatic heterocyclic group, optionally substituted by halo, alkyl, trifluoromethyl or trifluoromethoxy and a saturated or partially unsaturated 5 to 8 membered heterocyclic group containing one or more heteroatoms selected from nitrogen, oxygen or sulphur wherein the heterocyclic group is optionally substituted by one or more C₁₋₃alkyl groups, hydroxy, fluoro, benzyl or an amino group –NR^xR^y in which R^x and R^y independently represent H or C₁₋₄alkyl;

R³ represents a group of formula (CH₂)_nCOOR⁷

in which n is 0, 1, 2, 3 or 4 and R^7 represents a C_{4-12} alkyl group, a C_{3-12} cycloalkyl group or a $(C_{3-12}$ cycloalkyl) C_{1-3} alkyl—group each of which is optionally substituted by one or more of the following: a C_{1-6} alkyl group; fluoro, amino or hydroxy, or

BEST AVAILABLE COPY

WO 2004/111033

PCT/SE2004/000969

34

R⁷ represents a group -(CH₂)_aphenyl in which a is 0, 1, 2, 3 or 4 and the phenyl group is optionally substituted by one or more groups represented by Z which may be the same or different or

- R⁷ represents a saturated or partially unsaturated 5 to 8 membered heterocyclic group containing one or more of the of the following: oxygen, sulphur or nitrogen; wherein the heterocyclic group is optionally substituted by one or more C₁₋₃alkyl groups, C₁₋₃acyl groups, hydroxy, amino or benzyl; or
- R³ represents a group of formula -(CH₂)₀-O-(CH₂)_p- R⁸ in which o represents an integer 1, 2, 3 or 4 and p represents an integer 0, 1, 2, 3 or 4 and R⁸ represents a C₁₋₁₂alkyl group optionally substituted by one or more of the following: a C₁₋₆alkyl group; fluoro, hydroxy, or an amino group -NR^xR^y in which R^x and R^y independently represent H or C₁₋₄alkyl;
 - or R⁸ represents phenyl optionally independently substituted by one or more Z groups or R⁸ represents an aromatic heterocyclic group or a saturated or partially unsaturated 5 to 8 membered heterocyclic group containing one or more of one following: oxygen, sulphur or nitrogen wherein each of these rings is optionally substituted by one or more groups represented by Z which may be the same or different;
- R³ represents a group of formula -(CH₂)_qR⁹ in which q is 2, 3 or 4 and R⁹ represents a C₃.

 12cycloalkyl group, phenyl, an aromatic heterocyclic group or a saturated or partially unsaturated 5 to 8 membered heterocyclic group containing one or more of one following: oxygen, sulphur or nitrogen wherein each of these rings is optionally substituted by one or more groups represented by Z which may be the same or different; or

 R^3 represents a group of formula -(CH₂)_m-O-(CO)- R^{10} in which m represents an integer 0, 1, 2, 3 or 4, and in which R^{10} represents a C₁₋₁₂alkyl group optionally substituted by one or more fluoro, hydroxy, or amino or R^{10} represents a group of formula -(CH₂)_q R^9 in which q and R^9 are as previously described; or

30

35

R³ has the following formula:

 R^{11} represents hydroxy, fluoro, carboxy, a C_{1-6} alkoxycarbonyl group or an amino group - NR^xR^y in which R^x and R^y independently represent H or C_{1-4} alkyl;

d is 1, 2 or 3, and

R¹² represents H or a C₁₋₃alkyl group, or

R³ represents a group of formula CONH- R², in which R² is a piperidinyl ring substituted by a C₁₋₆alkanoyl group or R³ represents a group -COG in which G is a dihydroindole or a dihydroisoindole, linked through nitrogen to the carbonyl, and pharmaceutically acceptable salts thereof.

2. A compound according to claim 1, wherein R³ has the following formula:

15

10

 R^{11} represents hydroxy, fluoro, carboxy, a C_{1-6} alkoxycarbonyl group or an amino group - NR^xR^y in which R^x and R^y independently represent H or C_{1-4} alkyl; d is 1, 2 or 3,

 R^{12} represents H or a $C_{1\text{--}3}alkyl$ group, and pharmaceutically acceptable salts thereof.

20

3. A compound according to any of the preceding claims, wherein R^1 and R^2 each represent phenyl independently optionally substituted by one or more chloro.

BEST AVAILABLE COPY

WO 2004/111033

15

20

25

PCT/SE2004/000969

- 4. A compound according to any of the preceding claims, wherein R³ represents C₄₋₁₂alkoxycarbonyl.
- 5. A compound according to any of the preceding claims, wherein R³ represents a benzyloxymethyl group optionally substituted by Z in the phenyl ring of the benzyl group.
- 6. A compound according to any of the preceding claims, wherein R³ represents a group C(O)O-Het wherein Het is piperidino, morpholino or pyrrolidino.
- 7. A compound according to any of the preceding claims, wherein R¹ and R² each represent 4-chlorophenyl.
 - 8. A compound according to any of the preceding claims, wherein d is 1 and R^{11} is hydroxyl, amino or a $C_{1.6}$ alkoxycarbonyl group.
 - 9. A compound according to any of the preceding claims, wherein d is 2 and R¹¹ is F and both fluoros are attached to the same carbon on the cyclohexyl ring.
 - 10. A compound according to any of the preceding claims, wherein R¹² is H.
 - 11. A compound according to any of the preceding claims, wherein the aromatic heterocyclic group is furyl, pyrrolyl, thienyl, oxazolyl, isoxazolyl, imidazolyl, pyrazolyl, oxazolyl thiazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl or 1,3,5-triazenyl.
 - 12. A compound according to any of the preceding claims, wherein the aromatic heterocyclic group is pyrrolyl, thienyl, imidazolyl, oxazolyl or pyridyl.

WO 2004/111033

PCT/SE2004/000969

- 13. A compound according to any of the preceding claims, wherein the saturated or partially unsaturated 5 to 8 membered heterocyclic group is tetrahydrofuranyl, tetrahydropyranyl, pyrrolidinyl, morpholinyl, piperidinyl or piperazinyl,
- 14. A compound according to any of the preceding claims, wherein the saturated or partially unsaturated 5 to 8 membered heterocyclic group is tetrahydrofuran-3-yl, tetrahydropyran-4-yl, pyrrolidin-3-yl, morpholino, piperidino, piperidin-4-yl or piperazin-1-yl.
- 10 15. A compound selected from one or more of the following:
 - 5,6-bis(4-chlorophenyl)-N-(cis-2-hydroxypiperidin-1-yl)pyrazine-2-carboxamide,
 - 5,6-bis(4-chlorophenyl)-N-(trans-2-hydroxypiperidin-1-yl)pyrazine-2-carboxamide,
 - 5.6-bis(4-chlorophenyl)-N-(4-hydroxypiperidin-1-yl)pyrazine-2-carboxamide,
 - 5,6-bis(4-chlorophenyl)-N-(4,4-difluorocyclohexyl)pyrazine-2-carboxamide,
- N-(1-acetylpiperidin-3-yl)-5,6-bis(4-chlorophenyl)pyrazine-2-carboxamide,

 Tert-butyl 5,6-bis(4-chlorophenyl)pyrazine-2-carboxylate,
 - 5,6-Bis (4-chlorophenyl)-pyrazine-2-yl]-(1,3-dihydro-isoindol-2-yl)-methanone,
 - 2,3-bis(4-chlorophenyl)-5-{[(4-fluorobenzyl)oxy]methyl}pyrazine,
 - 2,3- bis(4-chlorophenyl)-5-[(piperidine-1-yloxy)carbonyl]pyrazine, and
- 20 pharmaceutically acceptable salts thereof.
 - 16. A compound of formula I as claimed in any previous claim for use as a medicament.
- 17. A pharmaceutical formulation comprising a compound of formula I, as defined in any of the claims 1-15 and a pharmaceutically acceptable adjuvant, diluent or carrier.
 - 18. Use of a compound of formula I according to any of the claim 1-15 in the preparation of a medicament for the treatment or prophylaxis of obesity, psychiatric disorders such as psychotic disorders, schizophrenia and bipolar disorders, anxiety, anxio-depressive disorders, depression, cognitive disorders, memory disorders, obsessive-compulsive disorders, anorexia, bulimia, attention disorders, epilepsy, and related conditions, and

BEST AVAILABLE COPY

WO 2004/111033

PCT/SE2004/000969

38

neurological disorders such as dementia, neurological disorders, Parkinson's Disease, Huntington's Chorea and Alzheimer's Disease, immune, cardiovascular, reproductive and endocrine disorders, septic shock, diseases related to the respiratory and gastrointestinal systems, and extended abuse, addiction and/or relapse indications.

19. A method of treating obesity, psychiatric disorders, psychotic disorders, schizophrenia and bipolar disorders, anxiety, anxio-depressive disorders, depression, cognitive disorders, memory disorders, obsessive-compulsive disorders, anorexia, bulimia, attention disorders, epilepsy, and related conditions, neurological disorders, neurological disorders, Parkinson's Disease, Huntington's Chorea and Alzheimer's Disease, immune, cardiovascular, reproductive and endocrine disorders, septic shock, diseases related to the respiratory and gastrointestinal system, and extended abuse, addiction and/or relapse indications, comprising administering a pharmacologically effective amount of a compound of formula I according to any of the claims 1-15 to a patient in need thereof.

20. A compound as defined in any of the claims 1-15 for use in the treatment of obesity.